

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



Dx

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/08	A1	(11) International Publication Number: WO 96/37213 (43) International Publication Date: 28 November 1996 (28.11.96)
(21) International Application Number: PCT/SG96/00004 (22) International Filing Date: 22 May 1996 (22.05.96) (30) Priority Data: 9500519-5 25 May 1995 (25.05.95) SG (71) Applicant (for all designated States except US): NATIONAL UNIVERSITY OF SINGAPORE [SG/SG]; 10 Kent Ridge Crescent, Singapore 119260 (SG). (72) Inventor; and (75) Inventor/Applicant (for US only): SIM, Meng, Kwoon [SG/SG]; National University of Singapore, Faculty of Medicine, Dept. of Pharmacology, 10 Kent Ridge Crescent, Singapore 119260 (SG). (74) Agent: APPLIED RESEARCH CORPORATION; Kent Ridge, P.O. Box 1016, Singapore 911101 (SG).		(81) Designated States: JP, SG, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: THE USE OF DES-ASPARTATE-ANGIOTENSIN I AS AN ANTI-CARDIAC HYPERTROPHIC AGENT (57) Abstract The use of des-Aspartate-angiotensin I (Arg-Val-Tyr-Ile-His-Pro-Phe-His-Leu) as an anti-cardiac hypertrophic agent is described. The compound, given either intravenously or orally, prevented the development of experimentally-induced cardiac hypertrophy in rats. Its action was dose-dependent and the maximum anti-cardiac hypertrophic effect was obtained at a dose of (i) 180 mg/day when given intravenously, and (ii) 285 mg/day when given orally.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

THE USE OF DES-ASPARTATE-ANGIOTENSIN I AS AN ANTI-CARDIAC HYPERTROPHIC AGENT

TECHNICAL FIELD

This invention relates to an anti-cardiac hypertrophic agent.

5 BACKGROUND ART

The interest in des-Aspartate-angiotensin I, a nine amino acid angiotensin peptide (Arg-Val-Tyr-Ile-His-Pro-Phe-His-Leu), as a peptide of the renin-angiotensin system was first generated when Blair-West and colleagues (Blair-West et al, *J. Clin. Endocrinol. Metab.*, 32:575-578 (1971)) postulated a biosynthetic pathway for the production of
10 angiotensin III by enzymatic NH₂-terminal degradation of angiotensin I to the nonapeptide and sequential action of angiotensin converting enzyme on this nonapeptide to produce the heptapeptide. Since then des-Aspartate-angiotensin I has been shown to be an excellent substrate of plasma and pulmonary angiotensin converting enzyme (Tsai et al, *J. Med. Chem.*, 18:1180-1183 (1975)) and that its pressor and steroidogenic actions are dependent on
15 its conversion to angiotensin III (Campbell et al, *Endocrinology*, 100:46-50 (1977)). Recently, we found that homogenates of rat aorta and hypothalamus degrade exogenous angiotensin I to mainly des-Aspartate-angiotensin I instead of angiotensin II and the enzyme responsible for the degradation was a specific aminopeptidase that was not inhibited by amastatin, bestatin and EDTA (Sim, *Biochem. Pharmacol.*, 45:1524-1527 (1993); Sim et al,
20 *Blood Pressure*, 3:260-264 (1994) and Sim et al, *Biochem. Pharmacol.*, 48:1043-1046 (1994)). Des-Aspartate-angiotensin I has also been shown to attenuate the pressor action of angiotensin II and angiotensin III in the brain (Sim and Radhakrishnan, *Eur. J. Pharmacol.*, 257:R1-R3 (1994)). Peripherally, it is able to potentiate the contractile action of angiotensin II on the rabbit aortic ring but to attenuate the contractile action of angiotensin III in the same
25 tissue (Sim and Yuan, *Eur. J. Pharmacol.*, 287:175-178 (1995)). These recent findings of ours seem to indicate that des-Aspartate I is a functional peptide that may have undefined specific actions in ensuring the normal functioning of the cardiovascular system.

DISCLOSURE OF INVENTION

In the course of studying the cardiovascular pharmacology of des-Aspartate-angiotensin I, the nonapeptide has been found to attenuate significantly the experimentally-induced cardiac hypertrophy in rat. It has been surprisingly discovered that des-

5 Aspartate-angiotensin I is effective in accordance with the present invention at an exceeding low dose, i.e. an i.v. dose of 180 ng (given over a period of 4 hours) per day for four days attenuates significantly the experimentally-induced cardiac hypertrophy in rats. Another significant finding is that, despite being a peptide, des-Aspartate-angiotensin I is equally effective in attenuating the cardiac hypertrophy when given orally at 285 mg per day for four
10 days. These findings show that des-Aspartate-angiotensin I is a highly specific anti-cardiac hypertrophic agent and is effective at concentrations that produce minimum or no secondary effects.

Therefore, the present invention is directed to the use of des-Aspartate-angiotensin I as an anti-cardiac hypertrophic agent or in the preparation of an anti-cardiac agent, for either
15 the prevention or treatment of cardiac hypertrophy or a pharmaceutical composition for preventing or treating cardiac hypertrophy comprising an effective amount of des-Aspartate-angiotensin I and a pharmaceutically acceptable carrier or diluent or a method for preventing or treating cardiac hypertrophy, which comprises administering to a subject in need of treatment an effective amount of des-Aspartate-angiotensin I or a packaged pharmaceutical
20 composition for preventing or treating cardiac hypertrophy comprising a container suitable for storing a pharmaceutical preparation, an effective amount of des-Aspartate-angiotensin I in said container, and instructions associated with said container giving instructions for the use of said des-aspartate-angiotensin I for preventing or treating cardiac hypertrophy.

MODES FOR CARRYING OUT THE INVENTION

25 In the practice of the method of the present invention, an effective amount of des-Aspartate-angiotensin I or a derivative or salt thereof, or a pharmaceutical composition containing the same, as described below, is administered to a subject, such as a human patient, via any of the usual and acceptable methods known in the art, either singly or in combination with other pharmaceutical agents such as captopril or other angiotensin
30 converting enzyme inhibitors. The compound or composition can thus be administered orally, by suppository, or parenterally (e.g., intramuscularly, intravenously, subcutaneously

or intradermally), and in the form of either solid or liquid dosage including tablets, suspensions, or solutions, as is discussed in more detail below. The administration can be conducted in single dosage form with continuous therapy or in single dose therapy ad libitum.

5 Useful pharmaceutical carriers for the preparation of the pharmaceutical compositions hereof can be solids, liquids or mixtures thereof; thus, the compositions can take the form of tablets, pills, capsules, powders, enterically coated or other protected formulations, sustained release formulations, erodible formulations, implantable devices or components thereof, microsphere formulations, solutions, suspensions, elixirs, aerosols, and
10 the like.

Water, saline, aqueous dextrose, and glycols are preferred liquid carriers, particularly (when isotonic) for injectable solutions. The carrier can be selected from various oils including those of petroleum, animal, vegetable or synthetic origin, for example, peanut oil, soybean oil, mineral oil, sesame oil, and the like. Suitable pharmaceutical excipients include
15 starch, cellulose, talc, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium stearate, sodium stearate, glycerol monostearate, sodium chloride, dried skim milk, glycerol, propylene glycol, water, ethanol, and the like. The compositions may be subjected to conventional pharmaceutical expedients such as sterilization and may contain conventional pharmaceutical additives such as preservatives, stabilizing agents, wetting or
20 emulsifying agents, salts for adjusting osmotic pressure, buffers, and the like. Suitable pharmaceutical carriers and their formulations are described in Martin, "Remington's Pharmaceutical Sciences", 15th Ed.; Mack Publishing Co., Easton (1975); see, e.g., pp. 1405-1412 and pp. 1461-1487. Such compositions will, in general, contain an effective amount of the active compound together with a suitable amount of carrier so as to prepare
25 the proper dosage form for proper administration to the host.

In one preferred embodiment, the therapeutic methods of the present invention are practiced when the relief of symptoms is specifically required or perhaps imminent; in another preferred embodiment, the method hereof is effectively practiced as continuous or prophylactic treatment.

30 In the practice of the therapeutic methods of the invention, the particular dosage of pharmaceutical composition to be administered to the subject will depend on a variety of

considerations including the stage of the disease or condition, the severity thereof, the schedule of administration, the age and physical characteristics of the subject, and so forth. Proper dosages may be established using clinical approaches familiar to the medicinal arts.

Although the initial work was conducted in a rat experimental model, it is expected
5 that the invention can be utilized in various mammals including, but not limited to, mice, rabbits and humans.

Example

Sources of Materials: Des-Aspartate-angiotensin I was obtained from Bachem (Bubendorf, Switzerland). Des-Aspartate-angiotensin I can be prepared by techniques well
10 known in the art. Adult Sprague Dawley rats (250-300 g) were obtained from the Animal Centre, National University of Singapore.

Induction of Cardiac Hypertrophy: The abdominal aorta of each animal was co-
arcted to the size of a 23-gauge hypodermic needle with a silk thread according to the
method described by Everett et al (*Hypertension*, 23:587-592 (1994)). Briefly, each animal
15 was anaesthetized with pentobarbital (5 mg/100 g, i.p.). An incision was made in the ventral
abdominal wall to access the suprarenal portion of the abdominal aorta. This portion of the
abdominal aorta was dissected free and a blunt 23-gauge needle was placed adjacent to the
aorta. A ligature was placed around the blunt needle and the aorta. The blunt needle was
then removed, leaving the aorta constricted to the size of the needle. The resulting
20 coarctation resisted the normal flow of blood from the heart to the lower portion of the body
and placed an extra load on the heart. This extra load is believed to cause hypertrophy of the
heart, especially the left ventricle.

Administration of Des-Aspartate-Angiotensin I: Following the operation, each
animal was administered one of the various doses of des-Aspartate-angiotensin I (dissolved
25 in saline) per day for four days. The nonapeptide was administered either intravenously via a
femoral vein catheter which was implanted during the co-arctation operation or orally via a 1
ml syringe with a blunt needle. The intravenous administration was carried out using a
microinjector which delivered 10 ml of the peptide solution per hour for four hours. For oral
administration, the peptide was dissolved in 0.5 ml saline. Control animals with co-arcted

abdominal aorta were administered saline instead of the peptide solution. Sham animals were animals that underwent the same surgical operations but their aortae were not co-arcted.

Determination of Cardiac Hypertrophy: On the fourth day following the co-arctation of the abdominal aorta, the animal was again anaesthetized with pentobarbital and the carotid and femoral blood pressure were determined via a carotid artery catheter and a femoral artery catheter, respectively. Each catheter was connected to a Gould Statham (P23 ID) pressure transducer. The transducers were in turn connected to a MacLab Quad Bridge Amplifier coupled to a MacLab/8 Virtual Instrument System which displayed the mean arterial blood pressure in mm Hg. The difference in the two readings indicated the extent of co-arctation. The heart of each animal was then excised and the weight of the ventricles was determined. The index of the ventricle weight (in mg) over the body weight of the animal (in g) was taken as the index of hypertrophy. For sham-operated animals the index was around 2.5, for aorta-co-arcted animal the index was above 3.7.

Results

The results of the study are summarized in Table 1. Des-Aspartate-angiotensin I has been found to be an effective agent in preventing the development of experimentally-induced cardiac hypertrophy. The anti-hypertrophic action is dose-dependent and its maximum action is brought about by an i.v. dose of 180 ng/day for four days or an oral dose of 285 mg/day for four days.

Table 1 Effects of des-Aspartate-angiotensin I on cardiac hypertrophy in rats

Dose	Hypertrophy Index ¹	CBP (mm Hg)	FBP (mm Hg)	BP (mm Hg)
No Administration				
Sham animals	2.53 ± 0.06	129 ± 13	129 ± 13	0
Intravenous Administration				
Control animals	3.77 ± 0.06	151 ± 19	107 ± 22	44
23 ng (19 pmol)	3.72 ± 0.18	158 ± 17	106 ± 13	52
45 ng (38 pmol)	3.51 ± 0.09	159 ± 25	119 ± 18	40
90 ng (76 pmol)	3.47 ± 0.13	156 ± 15	124 ± 14	32
180 ng (152 pmol)	3.17 ± 0.14	142 ± 22	111 ± 25	31
Oral Administration				
Control animals	3.75 ± 0.06	159 ± 18	110 ± 20	49
64 µg (63.5 nmol)	3.40 ± 0.10	153 ± 10	89 ± 27	64
128 µg (125 nmol)	3.23 ± 0.12	163 ± 27	121 ± 30	42
285 µg (250 nmol)	2.93 ± 0.09	171 ± 24	119 ± 33	52
590 µg (500 nmol)	3.13 ± 0.22	153 ± 22	110 ± 19	43
1180 µg (1000 nmol)	3.34 ± 0.16	154 ± 29	107 ± 25	46
1770 µg (1500 nmol)	3.57 ± 0.17	165 ± 14	123 ± 28	42

Each value is a means ± SEM obtained from 6 individual animals. Sham animals were animals that underwent the surgical operation but not the co-arctation of the abdominal aorta. Control animals were animals that underwent coarctation of the abdominal aorta but were given saline instead of the peptide solution. ¹Hypertrophy Index = ventricle weight in mg/body weight in g. CBP = mean arterial blood pressure obtained from the carotid artery catheter, FBP = mean arterial blood pressure obtained from the femoral artery catheter, BP = CBP - FBP. *Significantly different from the control (p < 0.05, Student's t-test).

1 INDUSTRIAL APPLICABILITY

- 2 The industrial applicability of the invention is primarily in the medical or health care
3 industry as an anti-cardiac hypertrophic agent in either the prevention or treatment of cardiac
4 hypertrophy.

CLAIMS

1 1. Use of des-Aspartate-angiotensin I as an anti-cardiac hypertrophic agent in
2 either the prevention or treatment of cardiac hypertrophy.

1 2. A pharmaceutical composition for preventing or treating cardiac hypertrophy,
2 comprising:

3 an effective amount of des-Aspartate-angiotensin I; and
4 a pharmaceutically acceptable carrier or diluent.

1 3. A method for preventing or treating cardiac hypertrophy, which comprises:
2 administering to a subject in need of treatment an effective amount of des-Aspartate-
3 angiotensin I.

1 4. A packaged pharmaceutical composition for preventing or treating cardiac
2 hypertrophy, comprising:

3 a container suitable for storing a pharmaceutical preparation;
4 an effective amount of des-Aspartate-angiotensin I in said container; and
5 instructions associated with said container giving instructions for the use of said des-
6 Aspartate-angiotensin I for preventing or treating cardiac hypertrophy.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SG 96/00004

A. CLASSIFICATION OF SUBJECT MATTER

IPC⁶: A 61 K 38/08

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC⁶: A 61 K 38/08

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, CAS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ENDOCRINOLOGY, Vol.108, No.2, February 1981 (Baltimore, USA), C. GARCIA DEL RIO et al.: "des-Asp- Angiotensin I: Its Identification in Rat Blood and Confirmation as a Substrate for Converting Enzyme", pages 406-412; totality.	1
A	DE 39 26 606 A1 (HOECHST) 14 February 1991 (14.02.91), abstract.	1,2,4

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"Z" document member of the same patent family

Date of the actual completion of the international search

05 August 1996 (05.08.96)

Date of mailing of the international search report

23 August 1996 (23.08.96)

Name and mailing address of the ISA/ AT
AUSTRIAN PATENT OFFICE
Kohlmarkt 8-10
A-1014 Vienna
Facsimile No. 1/53424/535

Authorized officer

Wolf

Telephone No. 1/53424/133

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SG 96/00004

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 3
because they relate to subject matter not required to be searched by this Authority, namely:
Method for treatment the human body by therapy (see also Rule 39.1(iv)
of the Regulations under the PCT).
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such
an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/SG 96/00004

In Recherchenbericht angeführtes Patentdokument Patent document cited in search report Document de brevet cité dans le rapport de recherche	Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication
DE A1 3926606	14-02-91	AT E 94409	15-10-93
		AU A1 60920/90	14-02-91
		CA B2 631914	10-12-92
		CA A1 2033089	10-02-91
		CA A1 9003958	12-08-92
		CA B6 2776444	17-03-93
		CA A5 2970633	02-01-93
		CA C0 5900273	11-10-93
		CA A1 417473	20-03-91
		CA B1 417473	15-09-93
		CA T3 2055933	16-11-94
		CA A0 9039336	00-08-90
		CA A2 9049666	01-04-91
		CA E 2055004	01-04-91
		CA A0 2055008	01-04-91
		CA A1 9039337	01-04-91
		CA A2 9039337	01-04-91
		CA A0 9039337	01-04-91
		CA A1 2864443	05-03-91
		CA A 9449666	05-03-91
		CA A 5231086	27-07-93
		CA A 9006327	29-05-91

AN. - 97-020931 [02]
AP - WO96SG00004 960522 EP960914540 960522; WO96SG00004 960522; [Based on
WO9637213] WO96SG00004 960522; US970776026 970519; [Based on
WO9637213]
PR - SG950000519 950525
TI - Use of des-aspartate-angiotensin I - as anti-cardiac
hypertrophic agent in prevention or treatment of cardiac hypertrophy
- DES ASPARTATE ANGIOTENSIN ANTI CARDIAC HYPERTROPHY AGENT PREVENT
TREAT CARDIAC HYPERTROPHY
PA - (UYSI-N) UNIV SINGAPORE NAT

Continue: Y / N

? Y

PN - WO9637213 A1 961128 DW9702 A61K38/08 Eng 013pp
- EP0774972 A1 970528 DW9726 A61K38/08 Eng 000pp
- US5773415 A 980630 DW9833 A61K38/00 000pp
IC - A61K38/00 ; A61K38/08
CT - 1.Jnl.Ref; DE3926606
AB - WO9637213 The use is claimed of des-aspartate-angiotensin I
(Arg-Val-Tyr-Ile-His-Pro-Phe-His-Leu) (DAA) as an
anti-cardiac hypertrophic agent in the prevention or treatment of
cardiac hypertrophy.
- ADVANTAGE - DAA is a highly specific anti-cardiac hypertrophic agent
and is effective at concns. that produce minimal or no secondary
effects.

D6